



International Journal of Sports Science & Medicine

Research Article

The Effects of Aerobic Exercise at Hypoxic Condition during 6 Weeks on Body Composition, Blood Pressure, Arterial Stiffness, and Blood Lipid Level in Obese Women -

Hun-Young Park¹ and Kiwon Lim^{1,2*}

¹Physical Activity and Performance Institute (PAPI), Konkuk University, Seoul, Republic of Korea

²Department of Physical Education, Konkuk University, Seoul, Republic of Korea

***Address for Correspondence:** Kiwon Lim, Department of Physical Education, Konkuk University, Seoul, Republic of Korea, Tel: +82-2-450-3827; E-mail: exercise@konkuk.ac.kr

Submitted: 15 August 2017; **Approved:** 22 August 2017; **Published:** 22 August 2017

Citation this article: Park HY, Lim K. The Effects of Aerobic Exercise at Hypoxic Condition during 6 Weeks on Body Composition, Blood Pressure, Arterial Stiffness, and Blood Lipid Level in Obese Women. *Int J Sports Sci Med.* 2017;1(1): 001-005.

Copyright: © 2017 Lim K, et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



ABSTRACT

The purpose of this study was to determine the effectiveness of aerobic exercise at hypoxic condition during 6 weeks on body composition, blood pressure, arterial stiffness, and blood lipid level in 30 to 60 years old women. Thirty five obese women above 30 in body mass index and %body fat were volunteered to participate in the study as participants and were divided into three groups such as 20.9% O₂ normoxic training group (n = 12), 16.5% O₂ hypoxic training group (n = 11), and 14.5% O₂ training group (n = 12). The six weeks of training consisted 30 min of treadmill and cycle ergometer exercise, respectively (total 1 hour). Exercise intensity was maximal heart rate of 75% that developed Miyashita in 1985. Exercise frequency was 1 hour a day, 5 days a week, during 6 weeks. As a result, there was a significant decrease ($p < .05$) in body fat mass and percentage of body fat in all groups. The reduction rate of body fat mass and percent of body fat were larger in the both hypoxic training group compared to the normoxic training group. Therefore, body weight was significantly decreased ($p < .05$) only in 14.5% O₂ hypoxic training group. Both hypoxic training groups showed a greater decrease ($p < .05$) in blood pressure and pulse wave velocity than normoxic training group. In addition, total cholesterol and low-density lipoprotein cholesterol showed a greater decrease ($p < .05$) in both hypoxic training group compared with normoxic training group. Based on these results, our study demonstrated that 16.5% O₂ and 14.5% O₂ hypoxic training had a positive effect and possibility of efficiency on body composition, blood pressure, arterial stiffness and blood lipid level in obese women compared with normoxic training.

Keywords: Hypoxic training; Body composition; Blood pressure; Arterial stiffness; Blood lipid level; Obese women

INTRODUCTION

Obesity is characterized by an increase in adipose tissue due to a positive energy balance of dietary intake and energy consumption and is now recognized as a very important social problem in increasing morbidity and mortality in the public health field [1]. In particular, the increase in visceral fat cells by obesity reduces the vasodilation of vascular endothelial cells and increases the risk of arteriosclerosis [2]. Many obese people often exhibit chronic inflammatory response to adipose tissue, which induces the onset and development of a variety of obesity-related diseases and increases oxidative stress in adipose tissue. In particular, obesity has been reported to cause metabolic dysfunction such as insulin resistance, type 2 diabetes and non-alcoholic fatty liver, and causes systemic inflammation, oxidative stress, and cardiovascular related complications [3].

In general, dietary and exercise combined treatment is the most commonly used to prevent obesity and various diseases [4]. However, this method has been reported to require long-term treatment for more than 12 weeks and does not inhibit the increased appetite with exercise, resulting in no consistent results via a relatively large number of positive and negative results [5]. In addition, pharmacotherapy and surgery can be used in the treatment of obesity, but these methods are not recommended except for severe obesity patients because of the various adverse effects that can adversely affect health [6].

In recent years, it has been recommended to hypoxic therapy (e.g. aerobic exercise in hypoxic condition) as a way to treat and prevent obesity and various lifestyle diseases by suppressing appetite and minimizing adverse effects [7]. In general, exercise in a high altitude environment or artificial hypoxic condition is well known to improve athletic performance [8]. In addition, exercise at hypoxic condition basically improves carbohydrate and lipid metabolism efficiently, it regulate oxygen transport capacity, process, carbohydrate metabolism, and satiety by increase the production of transcription factor hypoxia-inducible factor (HIF) -1 [9]. In particular, the hypoxic condition strongly induces weight loss through regulation of ghrelin, leptin, insulin, neuropeptide A, which increases fat metabolism and satiety [10-12]. Therefore, it is reported that exercise in the hypoxic condition is more effective in a shorter period of weight loss than exercise in the normoxic condition [7]. Exposure and exercise in a hypoxic condition leads to a decrease in arterial stiffness and an increase in blood flow to the skeletal muscle via vasodilation of the arteriole and production of nitric oxide (NO)[13]. These mechanisms

are thought to play an important role in the prevention and treatment of various cardiovascular diseases induced by obesity through normalizing blood pressure and blood vessel elasticity.

Therefore, using a randomized and controlled design, this study aimed to investigate the effects of 6 weeks of aerobic exercise at hypoxic condition corresponding to 16.5% and 14.5% O₂ compared with exercise in normoxic condition. We measured the effects on body composition, blood pressure, arterial stiffness, and blood lipid level in obese women.

MATERIALS AND METHODS

Participants

To determine the sample size, an a priori power analysis was performed with G-power for the body composition parameter (percent of body fat) based on previous research [14], indicating that a sample size of total 30 participants (10 participants per group) would be required to provide 80% power at an α -level of 0.05. We anticipated a more than 10% dropout rate and aimed for a starting population of 36.

Thirty-six middle-aged obese women (30 to 55 year), who were not taking any medication and with >30 BMI (by Broca's index) and $>30\%$ in percent body weight were selected as participants of this study. These women were housewives with low levels of activity who had not performed any kind of exercise over the last 6 months. Applicants' menopausal status was not assessed. The participants consented by signature, after sufficient explanation of the experiment and an understanding of the possible adverse effects, and were randomly assigned into a normoxic training group (n = 12), 16.5% O₂ hypoxic training group (n = 11), and 14.5% O₂ hypoxic training group (n = 12). Thirty-five of the participants completed the study ($>95\%$ compliance), thus, only their data were used in the analyses. Data from the remaining one subject was discarded due to medication (n=1). There were no significant differences in physical characteristics among groups before training (Table 1). All procedures followed were in accordance with the ethical standards of the responsible committee on human experimentation and with the Helsinki Declaration.

Study Design

Thirty-five middle-aged obese women in a normoxic training group (n = 12), 16.5% O₂ hypoxic training group (n = 11), and 14.5% O₂ hypoxic training group (n = 12) performed training

**Table 1:** Selected Participants Characteristics (Mean \pm S.D.).

Variables	Normoxic training group	16.5% O ₂ hypoxic training group	14.5% O ₂ hypoxic training group
Environmental condition	Sea-level	2,000m simulated altitude	3,000m simulated altitude
Number	12	11	12
Age (yrs)	47.2 \pm 6.3	42.0 \pm 4.4	46.6 \pm 5.7
Height (cm)	157.7 \pm 3.9	157.6 \pm 6.3	158.3 \pm 5.3
Weight (kg)	65.6 \pm 7.4	62.8 \pm 8.9	64.6 \pm 9.7
Body fat mass(kg)	34.3 \pm 3.3	34.8 \pm 4.6	32.8 \pm 2.2

Note:S.D: Standard Deviation

session equivalent to 75% of maximal heart rate (HR_{max}) at each environment during 6 weeks. Training session consisted of treadmill (30 min) and bicycle (30 min) exercise at a total duration of 1 hour, 5 days per week, for 6 weeks. The HR_{max} was determined using the predicted formula (Female = 205 - 0.75 \times age). During treadmill exercise, all participants were monitored for heart rate (HR) using a HR monitor (Polar S610i, Finland) to maintain their target HR of 75% HR_{max} as much as possible, and exercise was performed for 30 minutes while controlling the speed of the treadmill (Taeha IT-6025, Korea). For cycle ergometer exercise, all obese women entered their gender, age, weight, 75% HR_{max} into a cycle ergometer (Combi 75XLII, Japan) and attached a heart rate sensor to the earlobe. Then, the exercise load of the bicycle ergometer was automatically adjusted according to the heart rate information transmitted through the heart rate sensor attached to the earlobe. In this study, we designed a study to measure the effectiveness of hypoxic training vs normoxic training in obese women. Therefore, we analyzed body composition (weight, free fat mass, body fat mass, and percent of body fat), blood pressure (systolic and diastolic), arterial stiffness (pulse wave velocity in the upper and lower limbs), and blood lipid level (free fatty acid; FFA, total cholesterol; TC, high-density lipoprotein cholesterol; HDL-C, and low-density lipoprotein cholesterol; LDL-C) before and after 6 weeks of training.

A nitrogen generator (Separation & Filter Energy Technology Cooperation, Korea) was used to simulate 16.5% O₂ and 14.5% O₂ hypoxic conditions. The various hypoxic conditions were simulated by introducing nitrogen into the environmental chamber (width 6.5 m \times length 7.5 m \times high 3 m), using a nitrogen generator with the capacity to simulate normobaric hypoxic conditions for altitudes of up to 6000 m. The temperature within the environmental chamber was maintained at 20 \pm 2°C and the humidity was maintained at 60 \pm 2% for all conditions.

Measurements

Height was measured as the distance between the bottom of the foot and top of the head using a stadiometer (PKS-1008, JAPAN).

Body weight, free fat mass, body fat mass, percent of body fat were measured using an X-SCANPLUS (Jawon medical, Korea). All participants fasted overnight prior to the measurement of their body composition. They wore light weight clothing and were asked to remove any metal items.

Blood pressure was measured using a mercury sphygmomanometer (Sankei, Japan). All participants were examined at the same time as possible and had a stabilization time of at least 10 minutes before the test. The measurement was performed twice and the mean value was used as the measurement value.

The arterial stiffness was evaluated by pulse wave velocity (PWV) and it was measured using PWC 3.0 (Korea Medical Technology, Korea), a noninvasive device. All subjects were placed in a sitting position, two electrodes and one electrode were attached to the left arm and the right arm, respectively, and then the PWV of the upper limb was measured after inserting the sensor into two thumbs. Then, the sensor attached to the thumb was replaced with a big toe, and the PWC of the lower extremity was measured again. The result is recorded the mean value by selecting stable interval for 20 seconds from the data measured for 30 seconds.

For the analysis of blood lipid level, about 4 mL of blood was collected in the median antebrachial vein after 10 hours of fasting before and after 6 weeks of training at each environmental condition. After that, it was put into a SST (serum separating tube), centrifuged at 3,500 rpm for 10 minutes, and serum was collected and stored at -70°C for quick freezing. Then, the Green Cross Medical Foundation (Certified organization in The Korea Society for Laboratory Medicine) analyzed blood samples; a modular analytic PE (Roche, Germany) will be used to evaluate blood lipid levels. The enzymatic colorimetric assay method was used by FFA kit (Roche, Germany), CHOL kit (Roche, Germany), HDL-C plus 3rd generation kit (Roche, Germany), and LDL-C plus 2nd generation kit (Roche, Germany).

Statistical Analysis

All statistical analyses were conducted using SPSS 23.0 (IBM Corp., Armonk, USA) for windows. Data are presented as the mean \pm SD. The Kolmogorov-Smirnov test was used to ensure that all data had a normal distribution. All data were presented as mean \pm S.D. Paired *t* tests and analysis of variance (ANOVA) were used to evaluate within-participant differences between environmental conditions. The level of significance was set a priori at 0.05.

RESULTS

Body composition data (body weight, free fat mass, body fat mass, and percent of body fat), measured before and after training, is reported in Table 2. There was a significant decrease ($p < .05$) in body fat mass and percentage of body fat in all groups. The reduction rate of body fat mass and percent of body fat were larger in the both hypoxic training group compared to the normoxic training group. Therefore, body weight was significantly decreased ($p < .05$) only in 14.5% O₂ hypoxic training group.

Blood pressure (systolic and diastolic) data before and after training is shown in table 2. Systolic blood pressure was significantly decrease ($p < .05$) in the 16.5% and 14.5% O₂ hypoxic training group by 6 weeks of training. Diastolic blood pressure decreased significantly ($p < .05$) in all groups, but both hypoxic training group showed a larger decrease tendency than the normoxic training group.

As shown in table 2, arterial stiffness was evaluated by measuring the PWV at left hand, right hand, left foot, and right foot. The PWV showed no change by 6 weeks of normoxic training. However, the 16.5% O₂ hypoxic training group showed a significant decrease ($p < .05$) in PWV at left hand, right hand, and left foot. The 14.5% O₂ hypoxic training group showed improved arterial stiffness through a significant reduction ($p < .05$) in PWV at all sites.

Table 2 depicts before and after training data for FFA, TC, HDL-C, and LDL-C in all training groups. There was no significant difference in FFA and HDL-C by 6 weeks of training in all groups. TC significantly decreased ($p < .05$) in all groups. However, LDL-C showed a significant decrease ($p < .05$) only in the hypoxic group.

Table 2: Before- and after-training data for all outcomes by 6 weeks of training at each environmental condition (Mean ± S.D.)

Variables		Normoxic training group		16.5% hypoxic training group		14.5% hypoxic training group	
		Before	After	Before	After	Before	After
Bodycomposition	Weight (kg)	65.6 ± 7.4	65.1 ± 7.7	62.8 ± 8.9	62.3 ± 8.4	64.6 ± 9.7	63.4 ± 9.4*
	Free fat mass (kg)	40.5 ± 3.7	40.6 ± 3.8	38.4 ± 3.9	39.4 ± 3.7	40.8 ± 5.51	40.7 ± 5.20
	Body fat mass (kg)	22.6 ± 4.4	21.9 ± 4.9*	22.1 ± 5.8	20.6 ± 5.2*	21.3 ± 4.3	20.2 ± 4.4*
	Body fat (%)	34.3 ± 3.4	33.5 ± 3.6*	34.8 ± 4.6	32.7 ± 4.0*	32.8 ± 2.2	31.7 ± 2.7*
Bloodpressure	Systolic blood pressure (mmHg)	116.2 ± 10.5	113.1 ± 5.2	123.3 ± 11.6	116.5 ± 11.2*	121.8 ± 13.6	111.3 ± 11.5*
	Diastolic blood pressure (mmHg)	82.7 ± 9.1	79.0 ± 8.7*	76.7 ± 9.4	70.1 ± 7.3*	82.4 ± 8.9	73.6 ± 8.87*
Arterialstiffness(PWV)	Left hand (ms)	216.2 ± 14.7	208.2 ± 19.3	203.5 ± 13.7	195.4 ± 12.1*	211.6 ± 17.3	202.0 ± 25.2*
	Right hand (ms)	221.2 ± 23.8	222.3 ± 42.8	206.1 ± 11.3	197.9 ± 10.7*	218.2 ± 16.2	209.1 ± 15.0*
	Left foot (ms)	306.4 ± 25.1	291.5 ± 25.9	295.8 ± 17.9	289.6 ± 17.7*	308.6 ± 24.8	295.2 ± 26.5*
	Right foot (ms)	305.1 ± 20.3	291.1 ± 24.0	302.1 ± 18.2	300.1 ± 27.3	310.0 ± 24.8	297.4 ± 24.3*
Blood lipid level	Free fatty acid (uEq/L)	548.8 ± 169.7	516.8 ± 169.4	594.6 ± 181.7	465.2 ± 259.8	583.2 ± 165.0	533.8 ± 200.9
	Total cholesterol (mg/dL)	198.3 ± 33.2	175.7 ± 36.6*	186.6 ± 33.4	171.5 ± 42.5*	183.3 ± 13.7	172.6 ± 18.3*
	High density lipoprotein cholesterol (mg/dL)	50.7 ± 8.6	46.3 ± 9.6	47.4 ± 8.9	44.9 ± 8.8	47.5 ± 8.7	46.4 ± 6.6
	Low density lipoprotein cholesterol (mg/dL)	122.2 ± 41.4	102.4 ± 37.4	129.1 ± 33.2	102.1 ± 32.3*	120.7 ± 12.0	100.4 ± 15.1*

Note: S.D: Standard Deviation; PWV: Pulse Wave Velocity; * $p < .05$ vs. before in the same training group.

DISCUSSION

Many studies related to exercise training in high altitude or hypoxic conditions have been conducted primarily to improve aerobic exercise capacity of athletes [15]. Changes in substrate utilization and increased maximal oxygen consumption associated with improved aerobic exercise capacity are closely related to oxidative capacity of mitochondria in skeletal muscle [16]. In particular, improved aerobic exercise capacity is manifested by qualitative and quantitative adaptation of mitochondria [17]. Thus, aerobic exercise training in a hypoxic condition can be described as an effective treatment modality that can improve oxidative capacity and mitochondrial dysfunction in obese and type 2 diabetic patients [18,19]. However, based on these physiological mechanisms, studies on the application of aerobic exercise training at hypoxic condition for improving obesity and metabolic diseases are insufficient.

Therefore, Thirty-five middle-aged obese women in a normoxic training group (n = 12), 16.5% O₂ hypoxic training group (n = 11), and 14.5% O₂ hypoxic training group (n = 12) performed training session (treadmill and bicycle exercise at a total duration of 1 hour, 5 days per week, for 6 weeks) equivalent to 75% of maximal heart rate (HR_{max}) at each environment. As results, we found that 6 weeks of aerobic exercise training in the hypoxic conditions was more effective in improvement of body weight, systolic blood pressure, arterial stiffness, and LDL-C compared with the normoxic training. Therefore, we confirmed that hypoxic training could be applied as an effective treatment method for obesity and various lifestyle diseases.

In general, exercise training in the hypoxic condition increases the secretion of insulin, insulin like growth factor-1, erythropoietin, sex hormones such as androgens and testosterone, this increase in hormone secretion has been reported to increase muscle mass and increase basal metabolic rate [20-22]. In addition, Netzer, et al. [23]. Reported that exercise training in a hypoxic condition increases leptin secretion in adiposities, thereby increasing fat metabolism

and reducing body weight. Kayser & Verges [24], reported that the hypoxic condition stimulates the body to regulate various appetite-regulating substances related to weight control, and thus the body weight is reduced. Based on these various previous studies, the effect of improving the body composition in the hypoxic training group in this study seems reasonable.

Besides, exercise training in the hypoxic condition promotes the secretion of various metabolites such as NO, which induces relaxation of blood vessels in vascular endothelial cells, reduces the activity of the sympathetic nervous system, and increases blood flow and oxygen transport and utilization [7]. Wang, et al. [25]. Reported improved vasodilation of coronary and peripheral blood vessels due to increase NO and vascular endothelial growth factor production when sedentary subjects were given training in hypoxic environments. Nishiwaki, et al. [26]. Reported that intermittent moderate intensity training in a simulated 2,000 m altitude is more efficient in reducing atherosclerosis and increased flow mediated dilation. In this study, systolic blood pressure and arterial stiffness were improved in both the hypoxic training group compared with the normoxic training group, which was consistent with previous studies, and these results are valid.

Regarding blood lipid levels, Roberts, et al. [27] reported that exposure to the hypoxic condition increased the utilization of fat as a substrate, positively affecting blood lipid level. Tinkov & Aksenov [28]. Reported that 46 patients with ischemic heart disease and 45 men with coronary artery disease showed decrease of TC, LDL-C, and triglyceride after exposure and exercise in a hypoxic condition. Thus, they insisted that hypoxic condition is an important stimulus condition to improve blood lipid level and health of cardiovascular disease patients. Wiesner, et al. [19]. conducted a single blind study in overweight to obese subjects to test the hypothesis that training under hypoxia (HG, n = 24, FiO₂ = 15%) results in similar or even greater improvement in body weight and metabolic risk markers compared with exercise under normoxia (NG, n = 21, FiO₂ = 21%).



They conclude that in obese subjects, training in hypoxia elicits a similar or even better response in terms of physical fitness, metabolic risk markers, and body composition at a lower workload. The fact that workload and, therefore, mechanic strain can be reduced in hypoxia could be particularly beneficial in obese patients with orthopedic co morbidities. As with previous studies of this positive outcome, in this study, we also confirmed that exercise training in hypoxic condition is more effective in improving blood lipids such as LDL-C. The reduction of blood lipid levels in this study seems to be a complicated phenomenon appear to be caused by changes in dietary control mechanisms and improvement of body composition and cardiovascular function by exposure and training in hypoxic conditions. Therefore, various studies are needed to find out the precise mechanism and proper hypoxic stimulation condition.

CONCLUSION

The present study demonstrates that exercise training at 16.5% and 14.5%O₂ hypoxic conditions had a positive effect and possibility of efficiency on body composition, blood pressure, arterial stiffness and blood lipid level in obese women compared with normoxic training. It can be explained that there is a need to confirm the effect of hypoxic training on the basis of a review of various variables related to obesity and health during a longer treatment period in the future.

ACKNOWLEDGEMENT

All of the authors were involved in the conception, design, analysis, interpretation of results, and drafting of the article. The authors confirm that there are no conflicts of interest. This study was supported by a grant (NRF-2015M3C1B1019479) from the National Research Foundation funded by the Korean Government.

REFERENCES

- Drager LF, Li J, Reinke C, Bevans-Fonti S, Jun JC, Polotsky VY. Intermittent Hypoxia Exacerbates Metabolic Effects of Diet-Induced Obesity. *Obesity*. 2011; 19: 2167-2174. <https://goo.gl/yaqxri>
- Weil BR, Stauffer BL, Mestek ML, DeSouza CA. Influence of abdominal obesity on vascular endothelial function in overweight/obese adult men. *Obesity*. 2011; 19: 1742-1746. <https://goo.gl/Ljfu6c>
- Gade W, Schmit J, Collins M, Gade J. Beyond obesity: the diagnosis and pathophysiology of metabolic syndrome. *Clin Lab Sci*. 2010; 23: 51-61. <https://goo.gl/YtyU2d>
- Yoshizawa M, Maeda S, Miyaki A, Misono M, Saito Y, Tanabe K, et al. Effect of 12 weeks of moderate-intensity resistance training on arterial stiffness: a randomised controlled trial in women aged 32-59 years. *Br j sports med*. 2009; 43: 615-618. <https://goo.gl/NQ3DKF>
- Wu T, Gao X, Chen M, Van Dam RM. Long-term effectiveness of diet-plus-exercise interventions vs. diet-only interventions for weight loss: a meta-analysis. *Obes Rev*. 2009; 10: 313-323. <https://goo.gl/9emz4Z>
- Urdampilleta A, González-Muniesa P, Portillo MP, Martínez JA. Usefulness of combining intermittent hypoxia and physical exercise in the treatment of obesity. *J PhysiolBiochem*. 2012; 68: 289-304. <https://goo.gl/Meo9up>
- Millet GP, Debevec T, Brocherie F, Malatesta D, Girard O. Therapeutic Use of Exercising in Hypoxia: Promises and Limitations. *Front Physiol*. 2016; 7: 224. <https://goo.gl/fYqUqh>
- Park HY, Kim S, Nam SS. Four-week "living high training low" program enhances 3000-m and 5000-m time trials by improving energy metabolism during submaximal exercise in athletes. *J Exerc Nutrition Biochem*. 2017; 21: 1-6. <https://goo.gl/4Q4o5p>
- Vedam H, Phillips CL, Wang D, Barnes DJ, Hedner JA, Unger G, et al. Short-term hypoxia reduces arterial stiffness in healthy men. *Eur J Appl Physiol*. 2009; 105: 19-25. <https://goo.gl/Uy2VR3>
- Park HY, Nam SS, Kim SH, Kim MJ, Sunoo S. Effects of 10 weeks aerobic training in normobaric hypoxia on improvement of body composition, physical fitness, blood variables and vascular compliance. *Kor J Exer Nut*. 2010; 14: 7-16. <https://goo.gl/pPKjAv>
- Van Thienen R, Masschelein E, D'Hulst G, Thomis M, Hespel P. Twin resemblance in muscle HIF-1 α responses to hypoxia and exercise. *Front Physiol*. 2017; 7: 676. <https://goo.gl/MvfCve>
- Guerre-Millo M, Grosfeld A, Issad T. Leptin is a hypoxia-inducible gene. *Obes Res*. 2002; 10: 856-856. <https://goo.gl/CgAvhv>
- Wenger RH. Cellular adaptation to hypoxia: O₂-sensing protein hydroxylases, hypoxia-inducible transcription factors, and O₂-regulated gene expression. *FASEB J*. 2002; 16: 1151-1162. <https://goo.gl/Trm7GZ>
- Yingzhong Y, Droma Y, Rili G, Kubo K. Regulation of body weight by leptin, with special reference to hypoxia-induced regulation. *Int med*. 2006; 45: 941-946. <https://goo.gl/TK6JRZ>
- Park HY, Hwang H, Park J, Lee S, Lim K. The effects of altitude/hypoxic training on oxygen delivery capacity of the blood and aerobic exercise capacity in elite athletes - a meta-analysis. *J Exerc Nutrition Biochem*. 2016; 20: 15-22. <https://goo.gl/Yqk7cm>
- Blomstrand E, Rådegran G, Saltin B. Maximum rate of oxygen uptake by human skeletal muscle in relation to maximal activities of enzymes in the Krebs cycle. *J Physiol*. 1997; 501: 455-460. <https://goo.gl/7Y5v5D>
- Zoll J, Koulmann N, Bahi L, Ventura-Clapier R, Bigard AX. Quantitative and qualitative adaptation of skeletal muscle mitochondria to increased physical activity. *J Cell Physiol*. 2003; 194: 186-193. <https://goo.gl/Ho3u5E>
- Petersen KF, Dufour S, Befroy D, Garcia R, Shulman GI. Impaired mitochondrial activity in the insulin-resistant offspring of patients with type 2 diabetes. *N Engl J Med*. 2004; 350: 664-671. <https://goo.gl/r9sove>
- Wiesner S, Haufe S, Engeli S, Mutschler H, Haas U, Luft FC, et al. Influences of normobaric hypoxia training on physical fitness and metabolic risk markers in overweight to obese subjects. *Obesity*. 2010; 18: 116-120. <https://goo.gl/L4z2fV>
- Vogt M, Puntschart A, Geiser J, Zuleger C, Billeter R, Hoppeler H. Molecular adaptations in human skeletal muscle to endurance training under simulated hypoxic conditions. *J Appl Physiol*. 2001; 91: 173-182. <https://goo.gl/tZgrXW>
- Kon M, Ohiwa N, Honda A, Matsubayashi T, Ikeda T, Akimoto T, et al. Effects of systemic hypoxia on human muscular adaptations to resistance exercise training. *Physiol Rep*. 2015; 3:12267. <https://goo.gl/t6KX4X>
- Lundby C, Calbet JA, Robach P. The response of human skeletal muscle tissue to hypoxia. *Cell Mol Life Sci*. 2009; 66: 3615-3623. <https://goo.gl/qYKEVR>
- Netzer NC, Chytra R, Küpper T. Low intense physical exercise in normobaric hypoxia leads to more weight loss in obese people than low intense physical exercise in normobaric sham hypoxia. *Sleep Breath*. 2008; 12: 129-134. <https://goo.gl/Ri44s9>
- Kayser B, Verges S. Hypoxia, energy balance and obesity: from pathophysiological mechanisms to new treatment strategies. *Obes Rev*. 2013; 14: 579-592. <https://goo.gl/YK9yGP>
- Wang JS, Chen LY, Fu LL, Chen ML, Wong MK. Effects of moderate and severe intermittent hypoxia on vascular endothelial function and haemodynamic control in sedentary men. *Eur J Appl Physiol*. 2007; 100: 127-135. <https://goo.gl/b6sT68>
- Nishiwaki M, Kawakami R, Saito K, Tamaki H, Takekura H, Ogita F. Vascular adaptations to hypobaric hypoxic training in postmenopausal women. *The Journal of Physiological Sciences*. 2011; 61: 83-91. <https://goo.gl/ZDehYa>
- Roberts AC, Reeves JT, Butterfield GE, Mazzeo RS, Sutton JR, Wolfel EE, et al. Altitude and beta-blockade augment glucose utilization during submaximal exercise. *J Appl Physiol*. 1996; 80: 605-615. <https://goo.gl/qJDoV6>
- Tin'kov AN, Aksenov VA. Effects of intermittent hypobaric hypoxia on blood lipid concentrations in male coronary heart disease patients. *High Alt Med Biol*. 2002; 3: 277-282. <https://goo.gl/BXNBk8>